


4.1 Antiderivatives and Indefinite Integration



Anti Derivative
&
Uncle Function
are
my favourite
relations

- Antidifferentiation

Definition of Antiderivative

A function F is an **antiderivative** of f on an interval I when $F'(x) = f(x)$ for all x in I .

- Antidifferentiation by Trial and Error!

ex: Find a function, $f(x)$, given its derivative $f'(x)$.

a) $f'(x) = 2x$

ex: Find a function, $f(x)$, given its derivative $f'(x)$.

b) $f'(x) = 3x^2$

c) $f'(x) = x^4$

d) $f'(x) = -50 \sin x$

ex: Find a function, $f(x)$, given its derivative $f'(x)$.

e) $f'(x) = 2 \cos 2x$

f) $f'(x) = -x \sin x + \cos x$

Representation of Antiderivatives

If $\frac{d}{dx}[f(x)] = f'(x)$ then $f(x)$ is called the "general antiderivative" of $f'(x)$.

ex: Find the antiderivative: $f'(x) = 12x^7$

- Indefinite Integration

If $F(x)$ is any anti-derivative of $f(x)$ then the most general antiderivative of $f(x)$ is called an indefinite integral and denoted,

The diagram shows the formula $y = \int f(x) dx = F(x) + C$ with four labels in pink boxes connected by arrows:

- Variable of integration**: A box above the dx term with a downward arrow pointing to it.
- Constant of integration**: A box above the $+ C$ term with a downward arrow pointing to it.
- Integrand**: A box below the $f(x)$ term with an upward arrow pointing to it.
- An antiderivative of $f(x)$** : A box below the $F(x)$ term with an upward arrow pointing to it.

$$y = \int f(x) dx = F(x) + C.$$

++Differentiation and Integration are INVERSE Operations++

The inverse nature of integration and differentiation can be verified by substituting $F'(x)$ for $f(x)$ in the indefinite integration definition to obtain

$$\int F'(x) \, dx = F(x) + C.$$

Integration is the “inverse” of differentiation.

Moreover, if $\int f(x) \, dx = F(x) + C$, then

$$\frac{d}{dx} \left[\int f(x) \, dx \right] = f(x).$$

Differentiation is the “inverse” of integration.

These two equations allow you to obtain integration formulas directly from differentiation formulas, as shown in the following summary.

- Basic Rules

Differentiation Rules	Integration Rules
$\frac{d}{dx}[kx] =$	$\int k \, dx =$
$\frac{d}{dx}[kf'(x)] =$	$\int kf'(x) \, dx =$
$\frac{d}{dx}[f(x) \pm g(x)] =$	$\int [f(x) \pm g(x)] \, dx =$
$\frac{d}{dx}[x^n] =$	$\int x^n \, dx =$

*See printout.

Differentiation Rules	Integration Rules
$\frac{d}{dx}[\sin x] =$	$\int \sin x \, dx =$
$\frac{d}{dx}[\cos x] =$	$\int \cos x \, dx =$
$\frac{d}{dx}[\tan x] =$	$\int \sec^2 x \, dx =$
$\frac{d}{dx}[\csc x] =$	$\int \csc x \cot x \, dx =$
$\frac{d}{dx}[\sec x] =$	$\int \sec x \tan x \, dx =$
$\frac{d}{dx}[\cot x] =$	$\int \csc^2 x \, dx =$

Differentiation Rules	Integration Rules
$\frac{d}{dx}[e^x] =$	$\int e^x dx =$
$\frac{d}{dx}[a^x] =$	$\int a^x dx =$
$\frac{d}{dx}[\ln x] =$	$\int \frac{1}{x} dx =$

ex: Evaluate.

a) $\int x^5 dx =$

b) $\int 5\sec^2 x dx =$

c) $\int \sqrt{x} dx =$

ex: Evaluate.

d) $\int \frac{8}{x^2} dx =$

e) $\int \frac{8}{x} dx =$

f) $\int 2x dx =$

ex: Evaluate.

$$g) \int \frac{1}{(2x)^3} dx =$$

$$h) \int (4x^3 - 3^x + \sin x - 5e^x) dx =$$

$$i) \int \frac{x^7 - 5x^3 + 2x}{x^4} dx =$$

ex: Evaluate.

$$j) \int (1 + 3x) x^2 dx =$$

$$k) \int \frac{\sin x}{\cos^2 x} dx =$$

$$l) \int (1 + \cot^2 x) dx =$$

- Differential Equations

A differential equation is an equation involving a derivative.

- Differential Equations Have 2 Types of Solutions

1. General Solution - general antiderivative

2. Particular Solution - an antiderivative that passes through a given initial condition.

ex: $f'(x) = 3x^2 - 1$

a) Find the general solution.

b) Find the particular solution that satisfies the initial condition $f(2) = 4$.

- Total Distance (by hand)

ex: A particles moves on the x-axis so that its position at any time is given by: $x(t) = 4t^3 - 18t^2 + 15t - 1$

Find the total distance traveled by the particle from $t=0$ to $t=3$.

FR 2

A particle moves on the x -axis so that its velocity at any time $t \geq 0$ is given by $v(t) = 12t^2 - 36t + 15$. At $t = 1$, the particle is at the origin.

- (a) Find the position $x(t)$ of the particle at any time $t \geq 0$.
- (b) Find all values of t for which the particle is at rest.
- (c) Find the maximum velocity of the particle for $0 \leq t \leq 2$.
- (d) Find the total distance traveled by the particle from $t = 0$ to $t = 2$.

4.1-4.3 Extra Practice WKST

1.

If $f'(x) = 12x^2 - 6x + 1$, $f(1) = 5$, then $f(0)$ equals

(A) 2

(B) 3

(C) 4

(D) -1

(E) 0

4.1-4.3 Extra Practice WKST

2.

Find all functions g such that $g'(x) = \frac{5x^2 + 4x + 5}{\sqrt{x}}$

$$(A) \ g(x) = 2\sqrt{x}\left(x^2 + \frac{4}{3}x - 5\right) + C \quad (B) \ g(x) = 2\sqrt{x}\left(x^2 + \frac{4}{3}x + 5\right) + C$$

$$(C) \ g(x) = 2\sqrt{x}(5x^2 + 4x - 5) + C \quad (D) \ g(x) = \sqrt{x}\left(x^2 + \frac{4}{3}x + 5\right) + C$$

$$(E) \ g(x) = \sqrt{x}(5x^2 + 4x + 5) + C$$

4.1-4.3 Extra Practice WKST

3.

Determine $f(t)$ when $f''(t) = 2(3t + 1)$ and $f'(1) = 3$, $f(1) = 5$.

(A) $f(t) = 3t^3 - 2t^2 + 2t + 2$ (B) $f(t) = t^3 - 2t^2 + 2t + 4$

(C) $f(t) = 3t^3 + t^2 - 2t + 3$ (D) $f(t) = t^3 - t^2 + 2t + 3$

(E) $f(t) = t^3 + t^2 - 2t + 5$

4.1-4.3 Extra Practice WKST

4.

Consider the following functions:

I. $F_1(x) = \frac{\sin^2 x}{2}$

II. $F_2(x) = -\frac{\cos 2x}{4}$

III. $F_3(x) = -\frac{\cos^2 x}{2}$

Which are antiderivatives of $f(x) = \sin x \cos x$? (Hint: take the derivative of each and manipulate)

(A) II only (B) I only (C) I & III only (D) I, II, & III (E) I & II only

4.1-4.3 Extra Practice WKST

5.

A particle moves along the x -axis. The velocity of the particle at time t is $6t - t^2$. What is the total distance traveled by the particle from time $t = 0$ to $t = 3$?

- (A) 3 (B) 6 (C) 9 (D) 18 (E) 27

4.1-4.3 Extra Practice WKST

6.

A particle moves along the x -axis so that its acceleration at time t is $a(t) = 8 - 8t$ in units of feet and seconds. If the velocity of the particle at $t = 0$ is 12 ft/sec, how many seconds will it take for the particle to reach its furthest point to the right?

- (A) 6 seconds (B) 5 seconds (C) 3 seconds (D) 7 seconds (E) 4 seconds